


# AMERICAN METRIC BUREAU.

GENERAL OFFICES, 7 AND 8, SECOND STORY, 32 HAWLEY STREET, BOSTON.

CIRCULAR 22. — *The following article is issued separately for distribution as Circular 22, and will be sent to any physician applying. The author is the Chairman of the Bureau Committee on this subject. A brief circular, with equivalents, dangerous doses, etc., a new edition of the leaflet distributed last year, will soon be ready, and will be sent with this. This article is reprinted from the plates of the Boston Medical and Surgical Journal, and the old spelling, gramme, was then followed. Gram is now almost universally accepted as the correct spelling.*

## THE METRIC SYSTEM IN MEDICINE AND PHARMACY.<sup>1</sup>

BY T. B. CURTIS, M. D.



WE are all sufficiently familiar, in a general way, with the advantages claimed for the metric system of weights and measures. These, briefly enumerated, are: uniformity; simplicity; the decimal character; the fact that the various units have definite and readily apprehended relations to each other; a nomenclature which is self-defining and expressive of values; and, finally, security, the base being unalterable.

We also know the objections which have been raised against the proposed introduction of the metric system, and we have probably all been struck by the trivial and irrelevant character of several of them. The only objection which appears to me to possess any weight at all is one upon which our opponents habitually lay but little stress; yet it must be admitted that it has great force and deserves very careful consideration. I refer to the trouble and labor unavoidably attending any change of weights and measures. Routine, proceeding from traditions and habits of long standing and of numerous and complex ramifications, is firmly established. Is it worth our while to try to overcome the inertia of this routine? What are the inducements for us, whose main interest in the question is connected with the handling of drugs, to take a leading part in the proposed reform? In answer to this question I intend to pass in review, from the stand-point of the physician, certain advantages which can be claimed for the new system, and also to consider certain objections which have been brought forward by physicians against it.

First, it should be understood that the adoption of the metric system as used in Europe involves two separate and wholly independent innovations, consisting, primarily, in the substitution of a new system of weights and measures, with its nomenclature, for our present system; and, secondly, in the substitution of gravimetric methods for the system, partly gravimetric and partly volumetric, now in use.

With regard, in the first place, to the new system of weights and measures called the metric, this system recommends itself by several manifest merits. The advantage of uniformity with the usages estab-

<sup>1</sup> Read before the Suffolk District Medical Society, March 31, 1877.



lished among physicians in other countries is self-evident. In Germany and in France, which are, of all countries in which the English language is not spoken, those whose medical literature is most valuable to us, the metric system is used. In England weights and measures of the same names as ours are used, but they have different values; moreover, the English method of dealing with liquids is somewhat different from ours. All German, French, and English pharmaceutical formulas have therefore to undergo a process of translation, involving not only different weights and measures, but different methods of handling liquids, before they can be rendered available for use in this country. The adoption of the metric system, then, would bring us into agreement with the majority of civilized countries, while now we agree with none.

The metric system offers another advantage in that it provides denominations of weight applicable to the smallest quantities which the physician or the pharmacist can be called upon to prescribe or dispense. The advances achieved in chemistry and in therapeutics have provided a number of powerful drugs of which the doses are so minute as to oblige us to have recourse to very small fractions of our smallest unit of weight. The grain, which was suitable for dealing with the less powerful drugs formerly in use, is far too large a unit for the alkaloids and glucosides which modern chemistry has added to our *materia medica*. For handling these the centigramme and the milligramme of the metric system are eminently suitable.

Perhaps, however, the greatest advantage of the metric system accrues from the decimal character of its divisions. Some mathematicians have, I believe, asserted, on what to most of us must seem somewhat transcendental grounds, that the decimal system offers in itself no real advantage; that it has no *raison d'être* based upon any intrinsic properties of numbers or upon any mental attributes of mankind. This might possibly be true had other systems of numeration than those with which we are familiar originally been adopted. As a fact, however, our arithmetic is decimal. We count by series of tens, and higher and higher powers of ten constitute successive series of units. Hence the facility of decimal arithmetical operations. In the metric system, of which the divisions also proceed by series of tens, we change from one denomination to another, multiplying or dividing by ten, by one hundred, etc., by simply moving the decimal point to the right or to the left.

Another important advantage growing out of the arithmetical simplicity of a decimal division of our weights and measures is the readiness with which we should be enabled to appreciate quantitative ratios in our formulas and in the resulting pharmaceutical preparations. A few examples will suffice to make clear the difference in this respect between the old and the new system.



To illustrate, in the first place, the complexity and tediousness of the arithmetical processes involved in the use of our present method, and to show the difficulty of rapidly apprehending quantitative ratios, let us take the *mistura glycyrrhizæ composita*, or "brown mixture," of our Pharmacopœia. It contains the following ingredients: licorice, sugar, and gum arabic, of each  $\text{℥ss.}$ ; of camphorated tincture of opium,  $\text{f℥ij.}$ ; of wine of antimony,  $\text{f℥i.}$ ; of spirits of nitrous ether,  $\text{f℥ss.}$ ; of water,  $\text{℥fxxii.}$  The dose for an adult is a tablespoonful, for a child two years old a teaspoonful. Now, what are we giving with these doses? I once took the trouble to make the calculations, and found that a tablespoonful, equalling five drachms, contained: (1) of paregoric about  $\text{mxxxv.}$ , which by another similar calculation I found was equivalent to  $\text{gr. } \frac{1}{8}$  of opium; (2) of wine of antimony,  $\text{mxxviii.}$ , which yet another calculation showed to contain  $\text{gr. } \frac{1}{4}$  of tartar emetic. The other ingredients, being of no consequence medicinally, need not be estimated. Now these calculations, which are indispensable if we wish to know how much opium and how much tartar emetic our patient is getting with his repeated spoonfuls of brown mixture, are quite troublesome to make, and the results are difficult to remember. A similar task has to be performed in magistral prescriptions, when, instead of making out the value of the single dose of a complex and voluminous preparation, we proceed from the single dose with a view to formulating a mixture or pilular mass which shall be taken by repeated single doses, and last a stated time.

To illustrate the simplicity of metric formulas, on the other hand, let us take the *liquor hydrargyri bichloridi*, or *liqueur de Van Swieten* of the French codex. This solution contains: of the bichloride of mercury 1 gramme, or part, or unit of weight; of rectified spirit, 100 grammes, or parts; of water, 900 grammes, or parts. The liquid consequently contains one thousandth of its weight of the bichloride, and 5 grammes, the approximated value of a teaspoonful, contain 5 milligrammes of the drug.

Fowler's solution of arsenite of potassium, as formulated in the French codex, contains: of arsenious acid, 5 grammes; of carbonate of potassium, 5 grammes; of distilled water, 500 grammes; of alcohol, 15 grammes. After certain manipulations, comprising boiling, water is added to replace that lost, so that the whole weighs exactly 500 grammes, and thus contains one hundredth of its weight of arsenious acid. Of course the value in arsenious acid of any quantity by weight of the liquid is easily recognized.

One more example: let us take a magistral formula, the *sirop de Gibert*, or syrup of iodurated biniodide of mercury, a most valuable preparation. It contains: of the biniodide, 1 gramme; of iodide of potassium, 50 grammes; of water, 50 grammes, to which are added 2400



grammes of syrup of a certain specific gravity. The ratios involved are very simple, and we see at a glance, and remember with almost equal facility, that a tablespoon, holding 25 grammes of syrup, must contain one centigramme of the biniodide and 50 centigrammes of iodide of potassium. If now we should wish to formulate an analogous eight-ounce mixture we should find the necessary computations very tedious and difficult to perform accurately when in a hurry.

Our present system, then, is exceedingly complicated and difficult to manage in prescribing, in consequence of the absence of decimal divisions, and in consequence of the constant *changes of denomination* which result from processes of multiplication and division. One result, I believe, of this complexity of our system of weights and measures is the inordinate use in this country of *officinal*, in preference to *magistral*, or extemporaneous preparations. Many of our officinal remedies, to which we so largely resort, are complicated preparations, which have been elaborated at leisure by skilful and ingenious pharmacists, or by physicians of by-gone days, anterior to the time of rational therapeutics. These formulas, often embodying in one preparation such a number of active ingredients as to render it difficult or impossible to discriminate their effects upon the patient, are for the most part relics of an irrational and obsolete polypharmacy. Other analogous formulas are to be found scattered through text-books and treatises of theory and practice under the heading of "treatment." By many young physicians and students such formulas are copied out in memorandum-books, or learned by heart, and are used in a routine way, as if they represented simple drugs. Of course, when such habits of routine practice prevail, any advance in rational therapeutics is almost out of the question, and that self-education which should constantly be the chief object of our medication, after the benefit which we hope to confer upon the patient, is rendered difficult or impossible. Judging from the almost entire absence of such therapeutical formulas in text-books and treatises devoted to the instruction of French and German students and physicians, little or no assistance of this kind seems to be needed by them. If, now, magistral preparations were more extensively used, as would undoubtedly be the case if our system of weights and measures were easier to manage, then comparatively simple formulas would be readily improvised to meet the varying exigencies of individual cases, the tendency to polypharmacy would be diminished, the study and practice of rational therapeutics would be facilitated, and last, not least, the extensive use of patent medicines by physicians, which is the opprobrium of the medical profession of this country, would be lessened.

The second innovation comprised in the adoption by physicians of the system used in Continental Europe is the use of *gravimetric methods only*, for liquids as well as for solids. This is not an indispensable feat-



ure of the metric system, since the latter affords volumetric measures, decimally divided and having definite relations to the metric weights, which might with advantage be substituted for our present incongruous fluid measures. This innovation is, moreover, one which threatens to offer considerable difficulties and to meet with great opposition among physicians. If, then, it should appear likely to prove impracticable to bring about this change, we might for the present content ourselves with the simple substitution of metric weights *and measures* for those now in use.

But there can be little doubt that we shall in time generally recognize the intrinsic advantages of the gravimetric measurement of liquids, as well as the extrinsic advantage of conformity in this matter with European practice; and it may perhaps prove easier, if we are going to undertake any change in our system of weighing and measuring, to make it completely, once and for all.

The present state of the matter as regards the handling of liquids is as follows: while in England all solids are prescribed and dispensed by weight and all liquids by measurement, and while in all other countries solids and liquids alike are weighed, among us not only solids but also certain liquids, namely, acids, oils, honey, and chloroform, are ordered and dispensed by weight, other liquids being ordered by measurement.

What, now, will be the advantages and what the obstacles to the introduction of the gravimetric method for both solids and liquids? The first advantage to be looked for is conformity with the practice of all the nations which have adopted the metric system. The second and more important advantage lies in the greater accuracy in dispensing attainable by means of the gravimetric manipulation of liquids. Professor Maisch, after alluding to the frequent inaccuracy of the graduated measures now in use, to the difficulty of correctly reading the level of the liquid against the engraved scale, and to the additional uncertainties due to volatility and adhesion, says: "The greater convenience and correctness of weights have long since been recognized in the wholesale trade. Our Pharmacopœia even recognizes the correctness of this fact by having changed, in the last two revisions, all measures of the liquid acids, of chloroform, olive oil, and honey into weights. If these liquids are more conveniently and correctly handled by weight, why not likewise glycerine, syrups, tinctures, ethers, etc.?" He adds: "The dispensing of liquids by weight offers for all the reasons advanced by far greater accuracy than could be attained by measures."

Leaving, however, the facilitation of exact dispensing, which is a subject on which I am not qualified to express an opinion, I think there can be little doubt about the greater accuracy of prescriptions in which the quantities of all the ingredients, liquid as well as solid, are expressed by weight. A solid dissolved is a liquid, whose density, and conse-



quently whose exact bulk, we cannot readily ascertain. As Professor Maisch observes, solids dissolved in liquids add to the bulk of the latter. Moreover, certain liquids when mixed together contract, as is the case with alcohol or concentrated acids mixed with water. I think also that it is easier to recognize the ratio of a dissolved solid ingredient to its vehicle (whether the latter be water, syrup, or alcohol) when both are weighed than when the solids are weighed and the liquids measured, as in the *mistura glycyrrhizæ composita*, which was taken as an instance of our present way of dispensing.

The great difficulty which stands in the way of this innovation is that we have all contracted the habit of estimating liquids by measurement alone, and of remembering doses by bulk. We know only for distilled water the exact equivalence of our fluid measures in grammes; and unless we ascertain by experiment, or by calculation based on the knowledge of the specific gravity, the value in grammes of our present fluid measures (drachms and ounces) for every liquid, or at least for certain types of liquids, we shall have to treat all liquids by weight on the erroneous assumption that they have the same specific gravity as water.

The former solution of the difficulty is of course the correct one, and in due time we shall get to know all doses by weight, as we now know them by bulk. In furtherance of this end it is desirable that tables be constructed as soon as possible giving the value in grammes of the  $\text{f}\text{z}$  or  $\text{f}\text{z}$  of certain important liquid drugs (for example, *extract. ergot. fluid.*), as well as of certain types or categories of liquids, such as aqueous, alcoholic, and syrupy liquids. Future editions of the Dispensatory ought to give the corresponding values of every liquid by measurement and by weight in grammes.

But if, at present, instead of using metric measures of capacity for liquids, as has been proposed and is certainly feasible, we were to consider the fluid drachm of all liquids as equivalent to 3.69 grammes, this being the weight of one fluid drachm of distilled water, what would be the possible extent of the error involved? The inaccuracy of such a way of proceeding has been made a great "bugbear" by the opponents of the metric system. Lists of liquids of different densities have been gotten up, headed by ether, whose specific gravity is 0.728, and ending with chloroform, whose specific gravity is 1.490, water at 1.000, and syrups, from 1.310 to 1.320, coming between. The difference between the extreme specific gravities is as one to two, and a corresponding degree of inaccuracy in prescriptions has been predicted.

As a fact, however, neither ether nor chloroform is ever given in bulk or pure, unless it be by drops. Tinctures differ but slightly in specific gravity from water; aqueous solutions vary from 1.000 upwards, syrups at 1.320 representing the maximum specific gravity. The heaviest fluids, namely, syrups, rarely contain powerful drugs, and are hardly



ever given pure, being chiefly used for the purpose of imparting flavor. If, now, we should treat syrups as if of specific gravity 1.000, and assume the fluid drachm to weigh 3.69 grammes, the inaccuracy, as regards the amount of syrup alone, amounts to about twenty-five per cent. ; but in mixtures, into which syrups rarely enter in greater amount than one fifth of the whole, the inaccuracy would amount to four or five per cent. only of the whole mixture.

Now this inaccuracy, which could easily be reduced to a much more insignificant error, if not altogether annulled, by making a slight allowance for liquids heavier than water, has been held up as an insuperable objection to the substitution of the gravimetric method for the volumetric measurement of liquids. It seems to me, however, that those who urge this objection are claiming for their therapeutical operations a degree of accuracy which is rarely or never realized in practice.

It has been shown most clearly by Professor Maisch and by other authorities that the system actually in use involves a considerable degree of inaccuracy in the *prescription* and in the *dispensing* of medicines. There is a further and still greater margin of inaccuracy in the *administration* of medicines, owing to the false estimates in vogue with regard to the value of spoonfuls and drops, in which forms our drugs finally reach the patient.

A tablespoon is commonly supposed to contain half an ounce ; it really contains nearly six drachms, almost half as much again. A teaspoon, estimated to hold a fluid drachm, or sixty minims, really holds eighty minims, or one third as much again. A drop, supposed to be about equal to a minim, has values varying between a quarter of a minim and one minim and a quarter, the variation being as one to five ! The number of drops to a fluid drachm is with some fluids as low as forty-four, and with others as high as two hundred and seventy-six, if counted from a minim measure ; if counted from a bottle, on the other hand, some liquids give only forty-five drops to a fluid drachm, while others give as many as one hundred and eighty. For the same liquid the variation is often as two to three, according as the drops are counted from a bottle or from a minim measure. Thus a drachm of *tinctura opii* gives one hundred and forty-seven drops from a bottle and one hundred and six drops from a minim measure ; while a drachm of *tinctura ferri chloridi* gives one hundred and six drops from a bottle and one hundred and fifty-one drops from a minim measure. Yet, notwithstanding the great inaccuracy of this mode of measurement, drops are everywhere used in the prescription of the most concentrated and powerful liquid remedies, — such as, among others, tinct. digit., tinct. bellad., tinct. veratri viridis, tinct. aconiti radidis, acid. hydrocyan. dilut., liquor potassii arsenitis, etc. The danger of such grossly inaccurate measurements is obviated by the simple expedient, universally adopted, of giving in



every case small initial doses, which are gradually increased till the desired effect is attained, the final maximum dose in each case depending upon the quality of the drug and the susceptibility of the patient.

Thus we see that numerous causes of error beset our therapeutical operations in each of the necessary steps, as follows: (1.) *In prescriptions*, in consequence of the miscalculation of bulk when solids by weight and liquids by measurement are mixed. (2.) *In dispensing*, through the inaccuracy of graduates, the difficulty of reading correctly, volatility, adhesion, etc. (3.) *In administration*, in consequence of the prevailing erroneous estimates of spoonfuls and drops. I therefore venture to assert that the margin of inaccuracy is so great as to render insignificant a possible error of five per cent., due to the variations of specific gravity.

I believe, then, that the inaccuracy involved in neglecting, *for the present*, the differences of specific gravity of the various liquids would be more than compensated by the increase of accuracy which would accrue (after correction of our false estimates of the value of spoonfuls) from the substitution of the gravimetric handling of liquids for the mixed system, partly volumetric and partly gravimetric, now in use.

For all the reasons advanced I believe that the adoption by the medical profession of the metric system, as used in France, in Germany, and in many other countries, would be a most desirable consummation, and that it is expedient for us as physicians to make every effort to bring about this change.



THE FOLLOWING  
LECTURES

*Will appear shortly in the Boston Medical and  
Surgical Journal :—*

POST-PARTUM HÆMORRHAGE,

By Prof. R. A. F. PENROSE, of the University of Pennsylvania.

LACERATIONS OF THE CERVIX UTERI,

By Prof. T. GAILLARD THOMAS, of New York.

A series of four Lectures on the SPINAL CORD, by Prof. JOHN C. DALTON, of New York, entitled:—

1. Physiological Anatomy of the Spinal Cord and Nerves.
2. Transmission of Motor and Sensitive Impulses in the Spinal Cord and Nerves.
3. Crossed Action and Reflex Action of the Spinal Cord.
4. Reflex Action of the Cord (continued).  
Physiological Action of the Cord, as a Nervous Centre, during Health.

*Each number of the Journal now contains a lecture by some distinguished teacher. Specimen copies sent on application. For prospectus see next page.*



# THE BOSTON MEDICAL AND SURGICAL JOURNAL.

Established 1828.—Published Weekly.

THE PUBLISHERS OF THE BOSTON MEDICAL AND SURGICAL JOURNAL wish to call the attention of physicians to certain special features of this journal which make it exceptionally valuable as a medical periodical, and which commend it to the favor of professional gentlemen throughout the United States.

1. It is a weekly publication, thus possessing obvious advantages for the prompt presentation of medical intelligence, a great part of whose value lies in its freshness.

2. Being the only medical publication in New England, it possesses unusual opportunities for obtaining contributions to its pages, and is able to combine the special advantages of both a weekly and a monthly journal.

3. It contains *nothing* but original material. There are no borrowed articles, the current literature of the day being represented, in a manner peculiar to this journal, by a series of reports by experts in special departments of medical science, who give comprehensive and critical summaries of the most recent advances in their respective branches, and who constitute undoubtedly the strongest corps of the kind in the country.

4. It gives special attention to the department of reviews, which are fuller and more carefully prepared than in most medical journals.

5. Reports are received from a large number of medical societies, representing all departments of medicine. The facilities for obtaining clinical contributions from hospital practice have been greatly enlarged, embracing the most important hospitals of New England.

6. The correspondents' department has been organized with great care, and the chief medical centres of the world are here represented. A letter from one of these places appears in each issue of the JOURNAL.

7. A weekly table, showing the comparative mortality rates for the week, of thirteen large cities in the United States.

8. Editorially the JOURNAL aims to represent the profession of the whole country. It is one of the pioneers of medical science in the country, being, with one exception, the oldest now in existence. It was originated and has been maintained by no school or interest whatever beyond that of the science and practice of medicine and the medical profession.

The Editors and Publishers will spare no pains to keep the JOURNAL in the high and honorable position which it has so long occupied among American medical periodicals.

EDITOR.—DR. J. COLLINS WARREN.

EDITORIAL STAFF.—DR. A. L. MASON, DR. C. F. FOLSOM, DR. GEORGE B. SHATTUCK.

EDITORIAL REPORTERS.—DR. G. W. GAY, DR. F. W. GOSS, DR. HAMILTON OSGOOD.

DR. H. P. QUINCY, ANATOMICAL DRAUGHTSMAN.

## REPORTERS ON MEDICAL PROGRESS.

*Anatomy.*—THOMAS DWIGHT, M. D., late Professor of Anatomy of the Medical School of Maine.

*Chemistry.*—E. S. WOOD, M. D., Professor of Chemistry in Harvard University.

*Dentistry.*—THOMAS H. CHANDLER, D. M. D., Professor of Mechanical Dentistry, Harvard University.

*Dermatology.*—JAMES C. WHITE, M. D., Professor of Dermatology in Harvard University.

*Diseases of Children.*—D. H. HAYDEN, M. D., Physician to Out-Patients, Mass. General Hospital.

*Diseases of the Throat and Chest.*—FREDERIC I. KNIGHT, M. D., Instructor in Auscultation, Percussion, and Laryngoscopy, in Harvard University.

*Electro-Therapeutics.*—JAMES J. PUTNAM, M. D., Lecturer on Diseases of Nervous System in Harvard University.

*Gynecology.*—W. H. BAKER, M. D., Instructor in Gynecology in Harvard University.

*Materia Medica.*—ROBERT AMORY, M. D., Late Professor of Physiology in the Med. School of Maine.

*Mental Diseases.*—T. W. FISHER, M. D., Late Assistant Physician, Boston Insane Asylum.

*Military Surgery.*—GEORGE A. OTIS, M. D., U. S. A., Army Medical Museum, Washington, D. C.

*Obstetrics.*—W. L. RICHARDSON, M. D., Visiting Physician of the Boston Lying-in Hospital.

*Ophthalmology.*—O. F. WADSWORTH, M. D., Ophthalmic Surgeon, Boston City Hospital, and to Out-Patients Massachusetts General Hospital.

*Orthopedic Surgery.*—E. H. BRADFORD, M. D., Surgeon Carney Hospital.

*Otology.*—J. O. GREEN, JR., M. D., Aural Surgeon, Boston City Hospital.

*Pathology.*—R. H. FITZ, M. D., Assistant Professor of Pathological Anatomy in Harvard University.

*Physiology.*—HENRY P. BOWDITCH, M. D., Professor of Physiology in Harvard University.

*Public Hygiene and Forensic Medicine.*—FRANK W. DRAPER, M. D., Visiting Physician of the Boston City Hospital; Medical Examiner for Massachusetts.

*Surgery.*—J. COLLINS WARREN, M. D., Visiting Surgeon in the Massachusetts General Hospital; Instructor in Surgery at Harvard University.

*Syphilis.*—EDWARD WIGGLESWORTH, M. D., Lecturer on Syphilis, Harvard University.

*Theory and Practice of Medicine.*—DR. A. L. MASON, Physician to Out-Patients, Boston City Hospital.

*Genito-Urinary Surgery.*—T. B. CURTIS, M. D., Surgeon to Out-Patients in the Mass. Gen. Hospital.

**TERMS.**—Five dollars a year, payable in advance; single numbers, fifteen cents. Postage in all cases is paid by the Publishers. Communications for the JOURNAL should be addressed to the Editors, corner Beacon and Somerset streets, Boston; letters on business, to the Publishers, as below. Remittances may be made by draft, money-order, or registered letter to H. O. HOUGHTON & Co., Riverside Press, Cambridge, Mass.

HOUGHTON, OSGOOD AND COMPANY, BOSTON;

The Riverside Press, Cambridge.



# A COMPEND OF DIAGNOSIS

IN

## PATHOLOGICAL ANATOMY.

With Directions for Making Post Mortem Examinations. By Dr. JOHANNES ORTH.  
Translated by G. K. SABINE, M. D., and F. C. SHATTUCK, M. D. Revised by  
R. H. FIRZ, M. D. In one volume, 8vo, cloth, \$3.50.

In issuing a translation of this recent German work, the publishers have been convinced of its great practical value to all medical students. The directions for making an autopsy are given with great thoroughness and precision, the means of recognizing the various changes which may be found on the different parts of the body are briefly stated in a systematic manner, and the medico-legal bearing of the results of the investigation are always carefully attended to. The description of the gross appearances is also supplemented by thoroughly practical directions in the use of the microscope, when important additional information becomes necessary for exact diagnosis.

Dr. Orth was formerly the assistant of Professor Rindfleisch, and at present holds a like position under Professor Virchow. The teaching which he offers is consequently that of the highest authorities in the very centre of modern investigation in this department of medicine.

The publishers are the more desirous of calling attention to the translation, as it contains, through the kindness of the author, such additional information and corrections as he has prepared for the second edition of the original work, hereafter to be published. They have received many letters from prominent physicians and surgeons, warmly commending the book, and among them the following:—

COLLEGE OF MEDICINE, SYRACUSE UNIVERSITY.

SYRACUSE, N. Y., December 5, 1877.

Messrs. H. O. HOUGHTON & Co. *Gentlemen*:—Every physician ought to know how to make a *post mortem* examination. Not because any of *his* patients will ever die of a disease which baffled his diagnostic skill, but because old Grimness may have caught a brother practitioner napping, and have given to one of his “cases” the fatal stroke in a mysterious manner; or because the coroner may require his services in a legal investigation. So every physician should carefully read the admirable work of Prof. Orth on “*Diagnosis in Pathological Anatomy*.” For this book teaches—what so few know—how to make a neat *sectio cadaveris*. Prepared by a pupil of Virchow, the greatest pathologist in the world, it is not only fascinating, but authoritative on all points which it discusses. Cleverly translated, the book is one of the best additions to the best recent medical literature.

I thank you for the copy sent me. I heartily commend it to my classes, and to my professional brethren here and everywhere.

Respectfully yours,

H. D. DIDAMA.

PHILADELPHIA, December 7, 1877.

Messrs. H. O. HOUGHTON & Co. *Gentlemen*:—I have delayed the acknowledgment of the handsome translation of Orth’s “*Diagnosis in Pathological Anatomy*,” which you so kindly sent me, until I had time to read it carefully. Its clear and brief, but comprehensive, descriptions of the down-right facts of pathology are almost unsurpassed, and although I am at present teaching normal histology, I shall have and use many opportunities of recommending this work to our students.

Respectfully yours, etc.,

JOS. G. RICHARDSON.

OAKLAND, CAL., December 28, 1877.

Messrs. H. O. HOUGHTON & Co. *Gentlemen*:—Dr. Orth’s *Compend of Diagnosis in Pathological Anatomy* is a most admirably condensed and well arranged work—just the thing for students and busy practitioners. I shall take pleasure in recommending it to my class.

Yours truly,

J. K. WYTHE,

*Professor of Microscopy and Histology, Medical College of the Pacific.*

\*\*\* For sale by Booksellers, and by the Publishers,

HOUGHTON, OSGOOD AND COMPANY, BOSTON;

The Riverside Press, Cambridge.



# MEDICAL WORKS

**A Compend of Diagnosis in Pathological Anatomy.** By JOHANNES ORTH. Translated by G. K. SABINE, M. D., and F. C. SHATTUCK, M. D. Revised by R. H. FITZ, M. D. 8vo. Cloth, \$3.50.

**A Manual for Medical Officers of the Militia of the United States.** With a Complete Index. By EDWARD JACOB FORSTER, M. D. 12mo. Cloth, \$2.00.

**Civil Malpractice.** A Treatise on Surgical Jurisprudence, with Chapters on Skill in Diagnosis and Treatment, Prognosis in Fractures, and on Negligence. By M. A. McCLELLAND, M. D. 8vo. Cloth, \$3.50; sheep, \$4.50.

Your book is of the greatest value to the profession, and I am certain will become a standard authority. — LEWIS A. SAYRE, M. D., *Professor of Orthopedic Surgery, Fractures and Dislocations, and Clinical Surgery, Bellevue Hospital Medical College, New York*

**Gynecological Transactions.** Volume I. Being the Transactions of the American Gynecological Society at its First Annual Meeting, held in New York, September 13, 14, 15, 1876. 8vo. Cloth, gilt top, \$5.00.

As contributions to advanced gynecology they exceed in value anything which we have ever before seen collected in any one volume. Without exception, every paper has an interest and value attached to it which is intrinsic, and which reflects the best efforts of the respective authors. — *N. Y. Medical Record*

**The Cure of Rupture,** Reducible and Irreducible; also, of Varicocele. By GEORGE HEATON, M. D. Edited by J. HENRY DAVENPORT, M. D. 16mo. Cloth, \$1.50.

**Public Health, I., II.** Containing the Reports and Papers presented at the Annual Meetings of the American Public Health Association in 1873, 1874-1875. In two octavo volumes, 550 pages each, filled with important and valuable papers on topics of vital interest, not only to the profession, but to the entire community. Copiously illustrated with maps, charts, etc. Price, \$6.50 per volume. Volume III, 1875-1876, \$4.00.

*The Boston Daily Advertiser*, in noticing the second volume, says: "It ought not to be thought that it is a work valuable only to those who are interested in sanitary conditions as a matter of public concern. On the contrary, much of what it contains is exceedingly useful to private citizens whose care extends only to the safety of their own homes, or is even limited to their individual health."

**The Medical Register for New England.** By FRANCIS H. BROWN, M. D. 16mo. Cloth, \$2.50. A handy little volume containing statistical and practical information concerning Medical, Dental, and Pharmaceutical Societies and Associations, Hospitals and Dispensaries, Charitable Asylums, Institutions, and Societies, Professional Institutions of Education, Laws and Ordinances relating to Medical Men, Lists of Nurses, Surgical Instrument Makers, Chemists, Microscope Dealers, etc., etc. Together with an Autobiographical Record of the Members of New England Medical Societies.

**The Anatomy of the Head.** By THOMAS DWIGHT, M. D., Professor of Anatomy in the Medical School of Maine. In one volume. 8vo. Illustrated. Cloth, \$2.50.

One of the most important features of this work is six full-page lithographs of frozen sections of the head, which have been drawn with great care from original specimens. This is a method of demonstration which has received but little attention as yet from English and American anatomists; and the book, while designed especially for students, will also be of value to the general practitioner.

**Autumnal Catarrh (HAY FEVER).** By MORRILL WYMAN, M. D. With Illustrative Maps and Tables. 8vo. Cloth, \$2.00.

It contains a vast deal of information as to the various phases of the disease, the varying character of its exciting causes, its effects upon sufferers of different temperaments, and, more important than all, the best known methods of prevention and cure. — *Boston Transcript*.

**A Practical Guide to the Study of the Diseases of the Eye.** Their Medical and Surgical Treatment. By HENRY W. WILLIAMS, A. M., M. D. Fourth edition, revised and enlarged. Crown 8vo. Cloth, \$3.00.

**Recent Advances in Ophthalmic Science.** The Boylston Prize Essay for 1865. By HENRY W. WILLIAMS, A. M., M. D. Crown 8vo. Paper, \$1.00.

**Our Eyes and How to Take Care of Them.** 16mo. Cloth, \$1.00.

Dr. Williams's reputation as an oculist insures the value of these works.

**Boston Medical and Surgical Journal.** Vols. 92-97 (1875-1877), 8vo, cloth, per volume, \$3.50. Annual subscription \$5.00. Issued weekly, making two volumes a year. With one exception the oldest medical journal in the United States.

## IN PRESS.

**Manuals of Surgery.** By STEPHEN SMITH, M. D. A series of compact and useful hand books in various branches of Surgery, to be copiously illustrated, and neatly printed and bound

\*\*\* Those books already published will be sent by mail, post-paid, on receipt of price by the Publishers,  
HOUGHTON, OSGOOD AND COMPANY, BOSTON;

The Riverside Press, Cambridge.